



Direct Hair Implantation Pubic Restoration

MEDICAL BACKGROUND

-Woman of 53 years old
-Lost local hair due to local therapy with ointment for treating skin lesion

OBJECTIVE

To restore the damaged area

METHODS

DHI Direct Hair Implantation

BACKGROUND

In those body areas, hair loss problems may appear due to genetic problems, alopecia areata, and local factors like trauma, burn, surgical operation, laser therapy or chronic traction (i.e. trichotillomania).

BEFORE



METHODS

Treatment

- Donor area is the occipital scalp.
- Surgical methodology: the design takes place first, then the calculation of the necessary hair follicles is done, followed by the last three phases of the procedure (local anesthesia, extraction and placement).
- At the recipient area, endodermal local anesthesia is done since regional anesthesia or nerve block are not possible.
- Average hair density is around 20-25 hairs per cm²
- Direction of hairs : pubis: downwards

DESIGN -LOCATION

Pubic area: a reverse triangle over the pubic fat

RESULTS

Over the area of the mount pubis



RESULTS 5 months after DHI



Growth of more than 90%



CONCLUSIONS

- Single and double follicle grafts mixed are used throughout in both chest and pubic restoration .
- Random placement of the grafts is done at the limits of the designed area.
- The patient's desires regarding placement density and design should always be taken under consideration.
- The patient should avoid too much smoking and caffeine consumption as well as exercise for the first 10 days post-op.
- Light, loose and clean clothing is suggested for the first week after the Direct Hair Implantation session.
- Usually systematic antibiotics are given for a period of 5-7 days. It is up to the certified specialist's discretion to modify the dosage of the prescribed prednisolone. Cortisone can be given in a smaller dosage than in the scalp hair restoration.
- In the first two days post-op, the restored area should be sprayed with ringers solution.
- Wash the restored area on the 3rd day post-op by using a neutral soap.